



# Nucleoredoxin guards against oxidative stress by protecting antioxidant enzymes

Sophie Kneeshaw<sup>a,1</sup>, Rumana Keyani<sup>a,b,1</sup>, Valérie Delorme-Hinoux<sup>c</sup>, Lisa Imrie<sup>a,d</sup>, Gary J. Loake<sup>a</sup>, Thierry Le Bihan<sup>a,d</sup>, Jean-Philippe Reichheld<sup>c</sup>, and Steven H. Spoel<sup>a,2</sup>

<sup>a</sup>School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3BF, United Kingdom; <sup>b</sup>Department of Biosciences, Commission on Science and Technology for Sustainable Development in the South Institute of Information Technology, Islamabad 44000, Pakistan; <sup>c</sup>Université Perpignan Via Domitia, CNRS, Laboratoire Génome et Développement des Plantes, F-66860, Perpignan, France; and <sup>d</sup>Centre for Synthetic and Systems Biology (SynthSys), School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3JD, United Kingdom

Edited by Jonathan D. G. Jones, The Sainsbury Laboratory, Norwich, United Kingdom, and approved June 26, 2017 (received for review February 27, 2017)

Cellular accumulation of reactive oxygen species (ROS) is associated with a wide range of developmental and stress responses. Although cells have evolved to use ROS as signaling molecules, their chemically reactive nature also poses a threat. Antioxidant systems are required to detoxify ROS and prevent cellular damage, but little is known about how these systems manage to function in hostile, ROS-rich environments. Here we show that during oxidative stress in plant cells, the pathogen-inducible oxidoreductase Nucleoredoxin 1 (NRX1) targets enzymes of major hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)-scavenging pathways, including catalases. Mutant *nrx1* plants displayed reduced catalase activity and were hypersensitive to oxidative stress. Remarkably, catalase was maintained in a reduced state by substrate-interaction with NRX1, a process necessary for its H<sub>2</sub>O<sub>2</sub>-scavenging activity. These data suggest that unexpectedly H<sub>2</sub>O<sub>2</sub>-scavenging enzymes experience oxidative distress in ROS-rich environments and require reductive protection from NRX1 for optimal activity.

Nucleoredoxin | Thioredoxin | catalase | oxidative stress | reactive oxygen species

In eukaryotic cells, production of reactive oxygen and nitrogen species (ROS/RNS) is associated with many different developmental processes and stress responses. ROS/RNS such as superoxide (O<sub>2</sub><sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), and nitric oxide (NO) are continuously produced throughout cellular compartments as a by-product of metabolic pathways and in case of H<sub>2</sub>O<sub>2</sub> during periods of stress by a class of cell membrane-associated NADPH-dependent oxidases known as respiratory burst oxidase homologs (RBOHs) (1, 2). The inherent reactivity of these molecules and their ability to traverse membranes pose a significant danger to all cellular compartments (3), with common events including membrane lipid peroxidation, DNA damage, and irreversible hyperoxidation of proteins leading to loss of function (4). Consequently, ROS/RNS play key roles in cellular senescence and aging and have been linked to diseases such as cancer and the onset of neurodegenerative conditions (2, 5). Despite the dangers posed by ROS/RNS, they also function as indispensable short- and long-range signaling molecules in numerous cellular responses. Thus, ROS/RNS homeostasis needs to be carefully controlled to support signaling while preventing cellular damage.

To protect themselves against damage caused by oxidative stress and to redirect the actions of ROS/RNS into signaling pathways, plants and animals have evolved a plethora of antioxidant enzymes. The redoxin superfamily of oxidoreductases reverses the oxidative state of proteins modified by ROS/RNS. Allocation of Thioredoxin (TRX) activity is particularly important to protect and repair cell signaling proteins from undesired cysteine oxidation (6, 7). In addition, peroxidase family enzymes are up-regulated during periods of oxidative stress and directly metabolize ROS (1, 8). Although some peroxidases such as catalases use a haem center for dismutation of H<sub>2</sub>O<sub>2</sub>, other non-haem-containing peroxidases, including Ascorbate Peroxidases (APX) and Peroxiredoxins (PRX), often use thiol-based mechanisms to

decompose H<sub>2</sub>O<sub>2</sub> (9, 10). Notably, dysregulation of peroxidase enzymes is associated with cellular accumulation of H<sub>2</sub>O<sub>2</sub> and redox imbalance, resulting in accelerated aging and various pathological disorders (11–14). Although these findings indicate that this class of antioxidant enzymes is indispensable for maintenance of cellular redox, how they do so without sustaining significant structural damage in the hostile oxidative environments in which they function remains unknown.

Although cooperation between peroxidase and redoxin antioxidant systems during oxidative stress is apparent, the extent of this cooperation beyond the requirement for reducing power is currently unclear. In this study, we present evidence that activity of the plant immune-inducible TRX superfamily member, Nucleoredoxin 1 (NRX1), is necessary for the integrity of antioxidant systems. Our data unexpectedly suggest that H<sub>2</sub>O<sub>2</sub>-scavenging enzymes such as catalase suffer oxidative suppression in hostile ROS-rich environments and require oxidoreductase activity of NRX1 for optimal function.

## Results

A Pathogen-Inducible NRX Regulates Plant Immune Responses. Pathogen infection and subsequent activation of plant immune responses is associated with an oxidative burst and expression of selected TRX gene family members. Infection of wild-type (WT)

### Significance

Cellular accumulation of reactive oxygen species (ROS) such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) is associated with stress responses as well as aging. The reactive nature of ROS marks these molecules as a serious threat to cell integrity. Consequently, eukaryotic cells deploy numerous antioxidant enzymes that detoxify ROS to protect them from ROS-induced damage to proteins. Although the importance of antioxidant enzymes is well understood, how these proteins avoid becoming damaged in the hostile, ROS-rich environments in which they function remains unknown. We show that in plant cells the oxidoreductase Nucleoredoxin 1 (NRX1) protects antioxidant enzymes such as catalase from ROS-induced oxidation. Importantly, this protective effect of NRX1 boosted the H<sub>2</sub>O<sub>2</sub> detoxification capacity of catalase, thereby protecting the plant cell from oxidative stress.

Author contributions: S.K., G.J.L., T.L.B., J.-P.R., and S.H.S. designed research; S.K., R.K., V.D.-H., and L.I. performed research; S.K., R.K., V.D.-H., L.I., T.L.B., J.-P.R., and S.H.S. analyzed data; and S.K., T.L.B., J.-P.R., and S.H.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Data deposition: Raw and processed proteomic data were uploaded to ProteoSAFe, [massive.ucsd.edu/ProteoSAFe/](https://massive.ucsd.edu/ProteoSAFe/) (massIVE accession no. MSV000080429 and proteomeXchange accession no. PXD005591).

<sup>1</sup>S.K. and R.K. contributed equally to this work.

<sup>2</sup>To whom correspondence should be addressed. Email: [steven.spoel@ed.ac.uk](mailto:steven.spoel@ed.ac.uk).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1703344114/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1703344114/-DCSupplemental).

plants with the bacterial leaf pathogen *Pseudomonas syringae* pv. *maculicola* (Psm) ES4326 resulted in strong gene expression of a member of the NRX subfamily of TRX enzymes (Fig. 1A). Psm ES4326 infection also increased abundance of mixed disulfides between NRX1 and its substrates (Fig. 1B), indicating its oxidoreductase activity was pathogen-inducible. To investigate if NRX1 plays a functional role in plant immunity, we examined knockout lines (Fig. 1A and C). Mutant *nrx1* plants showed constitutive expression of pathogenesis-related (PR) genes but could be rescued by expression of Flag-tagged NRX1 from a constitutive 35S promoter (Fig. 1C and D). Elevated PR gene expression in *nrx1* was associated with enhanced disease resistance against Psm ES4326, which was comparable to WT plants immunized by the immune hormone salicylic acid (SA) or immunized by prior exposure to an avirulent pathogen (Fig. 1E and F). This autoimmune phenotype of *nrx1* mutants was dependent on SA, as mutation of the SA biosynthesis enzyme isochorismate synthase (ICS1) diminished immunity against Psm ES4326 (Fig. 1G).

Two Active Sites Make Differential Contributions to NRX1 Disulfide Reductase Activity. The NRX1 protein harbors three TRX-like domains, two of which contain an active site sequence consisting of WC(G/P)PC (Fig. 2A). Accordingly, NRX1 exhibits disulfide reductase activity (15), but the respective contributions of each active site have not yet been analyzed. In contrast to WT NRX1, mutation of either the first or third TRX-like domain in NRX1 (C55,58S) and NRX1(C375,378S) proteins, respectively, resulted in reduced disulfide reduction activity (Fig. 2B and Fig. S1). However, NRX1(C375,378S) was much more efficient than NRX1 (C55,58S), indicating that the majority of disulfide reduction activity occurred at the first TRX-like domain of NRX1. Simultaneous mutation of both the first and third TRX-like domains in the NRX1(C55,58,375,378S) protein completely abolished its ability to reduce insulin, demonstrating that these two domains contain all necessary active sites for disulfide reduction (Fig. 2B and Fig. S1). Because NRX1 proteins can dimerize (15), we investigated if active sites from different NRX1 proteins cooperate during disulfide reduction. Interestingly, when equal amounts of NRX1(C55,58S) and NRX1(C375,378S) were combined in an attempt to reconstitute the full enzymatic capacity of native NRX1 protein, we observed only a low level of activity that was comparable to NRX1 (C55,58S) protein alone (Fig. 2B). This suggests that the two active sites do not function independent from one another. Moreover, a single cysteine in one active site was not sufficient to support the second active site, as the two single cysteine mutants, NRX1(C58S) and NRX1(C378S), both exhibited reduced activity (Fig. 2C and Fig. S1), indicating a requirement for both active-site resolving cysteines for full oxidoreductase activity. Accordingly, disulfide reduction activity of the mutant proteins NRX1(C55,58,378) and NRX1(C58,375,378S) (Fig. S1), in which only a single cysteine of one active site was left intact, was completely abolished (Fig. 2C). Taken together, these findings imply that NRX1 reduces disulfide bonds of insulin by a dicysteine mechanism and that disulfide reduction may require cooperative or sequential interaction of a substrate with two different active sites of NRX1.

Identification of Stress-Induced Substrates of NRX1. Next we sought to identify NRX1 substrates using a previously designed capture strategy that exploits the mechanism by which disulfides are reduced by TRX family members (16, 17). Mutation of the second active-site cysteine of TRX enzymes prevents complete resolution of the disulfide reduction reaction, resulting in trapping of the substrate via an intermediate mixed disulfide bond. Therefore, we replaced the second cysteine with serine in each active site of NRX1. The resulting NRX1(C58,378S) mutant protein was unable to reduce insulin, confirming its inability to complete disulfide reduction reactions (Fig. 2D and Fig. S1). The NRX1 (C58,378S) protein was then immobilized on a column containing

NHS-activated resin and incubated with total protein extracted from Psm ES4326-inoculated plants. After trapping substrates via the formation of mixed disulfide bonds, the column was rigorously washed, substrates eluted with DTT, and finally identified by mass spectrometry (Fig. 2E). Two control columns were also included and treated identically. Nonspecific binding to the column was monitored by inclusion of a control column containing only




Fig. 1. NRX1 negatively regulates plant immune responses. (A) Plants were infected with Psm ES4326 ( $5 \times 10^5$  cells) for the indicated times. Expression of the NRX1 gene was analyzed and normalized against UBQ5. Error bars represent SD ( $n = 3$ ). (B) WT plants were infiltrated with or without Psm ES4326 ( $5 \times 10^5$  cells). Total protein was denatured and alkylated to capture true mixed disulfide intermediates between NRX1 and its substrates, separated by SDS/PAGE in the presence or absence of DTT, and analyzed by Western blot against NRX1. Indicated are free NRX1 monomer, mixed disulfide intermediates (NRX1-substrate), and total NRX1. (C) Expression of NRX1 was analyzed in WT, *nrx1-1*, *nrx1-2*, and 35S::Flag-NRX1 (in *nrx1-1*) plants. Gene expression was normalized against UBQ5 (error bars represent SD;  $n = 3$ ) and immunoblotting performed with anti-NRX1 and anti-HSP90 antibodies. (D) SA-dependent PR-1 and PR-2 gene expression. Error bars represent SD ( $n = 3$ ). (E) Plants were mock-treated or sprayed with 0.5 mM SA 24 h before infection with Psm ES4326 ( $5 \times 10^6$  cells). Growth of Psm ES4326 was assessed after 3 d. CfU, colony-forming units. Error bars represent statistical 95% confidence limits ( $n = 8$ ). Asterisks indicate statistically significant differences compared with the WT (Tukey-Kramer ANOVA test;  $\alpha = 0.05$ ,  $n = 8$ ). (F) Lower leaves were infiltrated with avirulent *P. syringae* pv. *tomato* DC3000/avrRpt2 ( $5 \times 10^7$ ) or 10 mM  $MgSO_4$ . After 2 d, upper leaves were infected with Psm ES4326 ( $5 \times 10^6$  cells) and pathogen growth assessed after 3 d. CfU, colony-forming units. Error bars represent statistical 95% confidence limits ( $n = 8$ ). Asterisks indicate statistically significant differences compared with the WT (Tukey-Kramer ANOVA test;  $\alpha = 0.05$ ,  $n = 8$ ). (G) Plants were infected with Psm ES4326 ( $5 \times 10^6$  cells) and pathogen growth assessed after 5 d. CfU, colony-forming units. Error bars represent statistical 95% confidence limits ( $n = 8$ ). \* $P < 0.05$  for statistical differences with WT and *nrx1*; \*\* $P < 0.05$  for statistical differences with WT, *ics1*, and *nrx1 ics1* (Tukey-Kramer ANOVA test;  $\alpha = 0.05$ ,  $n = 8$ ).




Fig. 2. Identification of substrates of NRX1 oxidoreductase activity. (A) Unique domain structures of AtNRX1 compared with AtTRXh5, a conventional immune-inducible TRX. (B–D) Oxidized insulin (130  $\mu$ M) was incubated with 0.3 mM DTT either alone (control) or together with 6  $\mu$ M native NRX1 or indicated NRX1 active-site mutants. Formation of reduced insulin was measured at 650 nm. Error bars indicate SD (n = 3). (E) Schematic of NRX1 substrate capture experiment, using immobilized mutant NRX1(C58,378S). (F) Venn diagram illustrating proteins identified in the substrate capture experiment. Each circle represents proteins found to be enriched ( $P < 0.05$ , ratio  $> 1.5$ ) in one of the following comparisons between columns: NRX1(C58,378S) column compared with WT NRX1 control column (pink); NRX1(C58,378S) column compared with resin-only control column (yellow); WT NRX1 control column compared with resin-only control column (blue). (G and H) GO term analysis was performed on the 74 targets enriched in the immobilized NRX1(C58,378S) mutant column for biological process (G) and molecular function (H) using Classification Supervisor on [bar.utoronto.ca/](http://bar.utoronto.ca/). Normalized frequencies, SD, and P values were determined from absolute values as described in ref. 36. Error bars represent SD.  $P < 0.05$  are printed bold.

quenched resin without any NRX1 protein. A second control column contained immobilized WT NRX1, allowing us to distinguish between targets that interacted as substrate or nonsubstrate. Compared with the WT NRX1 control column, 69 proteins were found to be enriched in the NRX1(C58,378S) column ( $P < 0.05$ , ratio  $> 1.5$ ) (Fig. 2F, pink circle). Additionally, 23 proteins were enriched in comparison with the resin-only control column ( $P < 0.05$ , ratio  $> 1.5$ ) (Fig. 2F, yellow circle). To further eliminate nonspecific background binding, a comparison was run between the WT NRX1 control column and the resin-only control column ( $P < 0.05$ , ratio  $> 1.5$ ). This identified 19 proteins, 7 of which overlapped with those previously identified as potential NRX1(C58,378S) substrates (Fig. 2F, blue circle) and were therefore excluded from further analysis. Thus, a total of 74 proteins were uniquely identified as potential substrates on the NRX1(C58,378S) column (Fig. 2F, pink and yellow circles excluding overlap with blue circle;

Dataset S1). A more stringent analysis still retained 45 potential NRX1 substrates with high confidence (Fig. S2A).

Gene ontology (GO) analysis of the identified targets revealed the most prominent biological processes were electron transport and energy pathways (Fig. 2G and Fig. S2B), consistent with the fact that these pathways often involve highly oxidative conditions. In agreement with our experimental approach, responses to stress as well as responses to abiotic or biotic stimuli were also found to be prominent biological processes (Fig. 2G and Fig. S2B). GO analyses for molecular function revealed that many potential NRX1 substrates exhibit structural molecule or enzymatic activity (Fig. 2H and Fig. S2C). As enzymes often contain active-site cysteines, these findings could suggest that NRX1 regulates their activity.

NRX1 Interacts with Enzymes of the Antioxidant  $H_2O_2$ -Scavenging Pathway. Upon further inspection of the entire list of identified potential NRX1 substrates, remarkably we identified many enzymes




Fig. 3. NRX1 protects plant cells from oxidative stress. (A) Diagram of the major  $H_2O_2$ -scavenging pathways. Enzymes enriched in the immobilized NRX1(C58, 378S) column are indicated in blue ovals and P values indicated above. (B) WT, 35S::Flag-NRX1 (Col-0), and 35S::Flag-NRX1 (cat2-2 cat3-1) plants were infected with Psm ES4326 ( $5 \times 10^5$  cells). Total protein was extracted under denaturing conditions and with the alkylating agent NEM to prevent nonspecific disulfide formation. Flag-NRX1 was immunoprecipitated and analyzed by Western blotting using an anti-CAT antibody. Anti-FLAG and anti-HSP90 were used as input controls. (C) Leaves were infiltrated with 5  $\mu$ M MV or ddH<sub>2</sub>O (control). Electrolyte leakage was measured as a percentage of total electrolytes. Error bars represent SD ( $n = 3$ ). Photographs were taken at 6 h. (D) Rosette leaf area (Left) and fresh weight (Right) were measured in 3-wk-old plants. Error bars represent SD ( $n = 20$ ). (E) Total leaf glutathione of leaves. Asterisks indicate a significant difference ( $P < 0.05$ ) between total glutathione levels of WT and *nrx1* plants. Reduced and oxidized glutathione levels are indicated by white and gray bars, respectively. The percentage reduction state of glutathione is indicated above bars. Error

of the main cellular  $H_2O_2$ -scavenging pathway (Dataset S1).  $H_2O_2$  is directly detoxified by both catalases and APX enzymes. Our quantitative mass spectrometry strategy identified as statistically significant several Arabidopsis catalases as well as the main cellular ascorbate peroxidase, APX1 (Fig. 3A, Fig. S3, and Dataset S1). Albeit outside the statistically significant range ( $P < 0.05$ ), we also consistently found the ascorbate regenerating enzymes, monodehydroascorbate reductase (MDAR1) and dihydroascorbate reductase (DHAR3), that underpin APX1 activity (Fig. 3A and Dataset S1). The pattern emerging from these data implicates NRX1 in the regulation of antioxidant enzymes.

Because of their prominent role in  $H_2O_2$  detoxification, we set out to validate the interaction of catalase enzymes with NRX1 in vivo. We performed a denaturing coimmunoprecipitation designed specifically to capture transient mixed disulfide interactions in vivo (see SI Materials and Methods) and found that catalase readily coimmunoprecipitated with Flag-tagged NRX1 in the WT but not in the *cat2 cat3* genetic background in which the two predominant catalase genes were knocked out (Fig. 3B) (18). These results confirm the mass spectrometry data and show that catalases are indeed substrates of NRX1 in vivo.

NRX1 Protects Cells Against Oxidative Stress. The identification of ROS-scavenging enzymes as targets of NRX1 prompted us to investigate if NRX1 plays a role in oxidative stress. We first assessed the cell death response of mutant *nrx1* plants treated with the ROS-generating chemical methyl viologen (MV). Compared with WT, mutant *nrx1* plants displayed faster and more pronounced MV-induced cell death than the WT (Fig. 3C). This is reminiscent of the ROS intolerant phenotype of mutant *cat2* plants (19), which, like *nrx1* plants also display autoimmune phenotypes (20). We considered the possibility that NRX1 and catalases may act within the same ROS detoxification pathway. To investigate this further, we studied the phenotypes of an *nrx1 cat2* double mutant. Mutation of NRX1 did not further aggravate the developmental phenotype of *cat2* mutants (Fig. 3D). We then assessed the cellular redox status of these mutants. Compared with WT, mutant *nrx1* plants accumulated significantly higher levels of total glutathione, but the ratio of oxidized versus reduced glutathione was normal (Fig. 3E). These data suggest that in absence of a stimulus, *nrx1* mutants experience low levels of oxidative stress that are managed by increased glutathione levels. Mutation of NRX1, however, was not able to aggravate further the redox status of *cat2* mutants, which as reported previously (19) contained very high levels of total and oxidized glutathione (Fig. 3E). These data suggest that NRX1 and CAT2 do not act additively to control the redox status of resting cells (Fig. 3E).

To assess if NRX1 and CAT2 function together or additively in stressed cells, we measured cell death in single and double mutants that had been treated with MV. As expected, both *nrx1* and *cat2* single mutants displayed increased levels of cell death in comparison with WT (Fig. 3F). However, the *nrx1 cat2* double mutant was as susceptible to oxidative stress as *cat2* single mutants, indicating that mutation of these genes also did not act additively during oxidative stress. Taken together, these data demonstrate that NRX1 is essential for protection against oxidative stress and suggest that it may act by modulating the activity of catalases.

Catalase Activity Is Controlled by NRX1-Mediated Reduction. Our findings suggest that catalases may be recipients of oxidative posttranslational modifications and may require NRX1 to remain in the reduced state. To examine this possibility, we performed a reductive switch assay in which specifically oxidized cysteines

bars represent SD ( $n = 3$ ). (F) Leaves were infiltrated with 5  $\mu$ M MV or ddH<sub>2</sub>O (control). Electrolyte leakage was measured as a percentage of total electrolytes. Error bars represent SD of the mean ( $n = 3$ ).

undergo DTT-dependent labeling with biotin followed by pull-down with streptavidin. Indeed, catalase was subject to oxidative modification, and regardless of the treatment, the level of oxidized catalase was greatly enhanced in *nrx1* mutants compared with WT, supporting the notion that NRX1 controls the oxidation state of catalase (Fig. 4A). To determine if NRX1 plays a role in modulating catalase activity, we examined catalase activity in protein extracts from untreated, MV-treated, and Psm ES4326-infected plants. Decomposition of  $H_2O_2$  could be largely attributed to catalase activity, as *cat2 cat3* double mutants displayed only residual activity (Fig. 4B and Fig. S4A). In striking contrast to WT, in all treatments *nrx1* mutants displayed significantly decreased catalase activity (Fig. 4B and Fig. S4A), despite comparable catalase protein levels between genotypes (Fig. S4B). A direct link between NRX1 activity and the ability of catalase to decompose  $H_2O_2$  was found by incubating plant extracts with recombinant WT NRX1 or mutant NRX1(C55,58,375,378S) protein. Addition of either recombinant protein to *cat2 cat3* extracts did not increase  $H_2O_2$  consumption, indicating NRX1 cannot detoxify  $H_2O_2$  by itself (Fig. 4C and Fig. S4C). By contrast, addition of recombinant NRX1 but not mutant NRX1(C55,58,375,378S) to *nrx1* extracts remarkably restored catalase activity to levels comparable to those found in WT extracts (Fig. 4C and Fig. S4C). Moreover, recombinant NRX1 significantly increased catalase activity in WT plant extracts, whereas mutant NRX1(C55,58,375,378S) protein was ineffective in this respect. In summary, these data demonstrate that NRX1 controls the oxidation status of catalase and thereby modulates its  $H_2O_2$  dismutation activity.

## Discussion

During periods of oxidative stress, ROS-scavenging by peroxidase enzymes is essential to maintaining cell integrity. However, how these enzymes withstand the toxic oxidizing conditions in which they function remains elusive. In this study, we demonstrate that in plants, NRX1, a member of the TRX superfamily of enzymes, maintains catalase enzymes in a reduced state, thereby protecting their  $H_2O_2$ -detoxifying activity and, importantly, ensuring efficient functioning of the antioxidant system during oxidative stress.

Using extracts from immune-induced plants, we identified 74 potential endogenous substrates of NRX1 (Fig. 2F). Although some of these are unlikely to be real substrates, strikingly we found that NRX1 engaged with enzymes of a major  $H_2O_2$ -scavenging pathway, including Arabidopsis catalases (Fig. 3A and Fig. S2). Indeed, we confirmed that catalase was a substrate of NRX1 *in vivo*, and epistasis experiments strongly suggested that NRX1 and the major leaf catalase CAT2 act together in the same  $H_2O_2$ -detoxification pathway (Fig. 3). The full extent to which catalase activity contributes to plant immune responses has only recently come to light. Catalases have previously been reported to bind SA, and a recent study showed that SA-induced suppression of CAT2 activity increased  $H_2O_2$  levels upon pathogen infection (21). Here we show that additionally, during oxidative stress and pathogen infection, CAT2 activity is regulated by NRX1. Whether this is a conserved role of NRX1 in other eukaryotes remains to be seen, but in addition to this study, NRX1 has been implicated in immune responses in both a crop plant and mammals (22, 23).

Catalases are widely reported to localize to the peroxisomes where they remove toxic  $H_2O_2$  generated in cellular metabolism. However, NRX1 lacks known peroxisomal targeting sequences (PTSs) and is localized to the cytoplasm and nucleus (15). Thus, it is unlikely that peroxisomal catalases are a substrate of NRX1. Nonetheless, using a coimmunoprecipitation trap, we found that NRX1 formed a mixed disulfide intermediate with catalase *in vivo* (Fig. 3D). In this respect, it should be noted that peroxisomal import of catalases is dependent on cytosolic interaction with the PTS import receptor, PEX5 (24, 25). PEX5 docks with the peroxisomal membrane where it releases its cargo into the peroxisome matrix. Under oxidizing conditions, however, PEX5 has been shown to undergo redox-dependent monoubiquitination at a




Fig. 4. NRX1 regulates redox modification and activity of catalase. (A) Protein extracts from untreated, MV- (5 M), or Psm ES4326- ( $5 \times 10^5$  cells) treated plants were alkylated and incubated with or without DTT (2 mM). Free thiols were labeled with biotin and pulled down with Streptavidin. Oxidized catalase (OX-CAT) was visualized by immunoblotting with an anticatalase antibody and shown relative to total catalase. (B) The ability of protein extracts from untreated, MV- (5 M), or Psm ES4326- ( $5 \times 10^5$  cells) treated plants to decompose  $H_2O_2$  (5 mM) was measured at  $A_{240}$  and leaf catalase activity ( $\mu\text{mol } H_2O_2 \cdot \text{mg} \cdot \text{protein}^{-1} \cdot \text{min}^{-1}$ ) calculated using the extinction coefficient of  $0.036 \text{ cm}^2 / \text{mol}$ . Error bars represent SE ( $n = 3$ ). \* $P < 0.05$  for statistical differences with WT and *cat2 cat3*; \*\* $P < 0.05$  for statistical differences with WT and *nrx1* (Student's t test). (C) As in B, but extracts were supplemented with 4 M recombinant NRX1 or NRX1(C55,58,375,378S) and 0.33 mM DTT. pNRX1, recombinant NRX1 protein added to reaction mixture; mut., mutant recombinant NRX1 protein. Error bars represent SE ( $n = 3$ ). \* $P < 0.05$  for statistical differences compared to mutant recombinant NRX1 protein (Student's t test).

cysteine residue in the N terminus, preventing it from being recycled back to the cytosol (26). Consequently, during periods of oxidative stress, PEX recycling is blocked, resulting in decreased peroxisomal import of PEX5 targets, including catalase (25). Accordingly, a recent report demonstrated that pathogen-mediated transient expression of CAT3 in tobacco leaves resulted in its localization to the peroxisome, cytoplasm, and plasma membrane (27). Our findings suggest that oxidation-induced retention of catalases in the cytoplasm may enable these critical detoxifying enzymes to become substrates of NRX1.

Catalases contain several surface-exposed cysteine residues that may be sensitive to oxidation (28–30). Therefore, it is surprising that these enzymes function efficiently in ROS-rich environments without incurring significant damage. Here we found that catalases are indeed subject to inhibitory oxidative modification and demonstrate that NRX1 may be dedicated to maintaining these enzymes in a reduced state to promote their H<sub>2</sub>O<sub>2</sub>-scavenging activity (Fig. 4). Studies on the green alga *Chlamydomonas reinhardtii* have demonstrated reversible, partial inactivation of catalase, which was attributed to redox regulation of a single cysteine residue that is conserved among *Arabidopsis* catalases (30). Oxidation of a single cysteine implies that rather than a disulfide, catalase may contain an S-nitrosothiol or sulfenic acid. Although the ability of NRX1 to reduce these modifications has not yet been demonstrated, they are targeted by conventional TRX family members (31–33). Regardless of these particulars, this study clearly demonstrates that NRX1-mediated reduction of catalase is an important mechanism for sustaining maximal catalase activity.

So does the advantageous effect of NRX1 on H<sub>2</sub>O<sub>2</sub> scavenging extend beyond regulation of catalases? Extraordinarily our proteomic capture experiments recovered nearly all major enzymes of the cytosolic glutathione/ascorbate-dependent H<sub>2</sub>O<sub>2</sub>-scavenging pathway, including APX, MDAR, and DHAR (Fig. 3 and Dataset S1), some of which have been identified individually in other proteomic screens for substrates of conventional TRX enzymes (see ref. 34 and references herein). The importance of the cytosolic glutathione/ascorbate cycle in H<sub>2</sub>O<sub>2</sub> scavenging is exemplified by reports that showed reduced levels of APX

protein or ascorbate deficiency resulted in high levels of oxidative stress, spontaneous cell death, and hyperresponsiveness to plant pathogens (11, 35). Given that cysteine oxidation appears prevalent in controlling activities of enzymes in this H<sub>2</sub>O<sub>2</sub>-scavenging pathway, it is highly likely that the interactions found here between NRX1 and these enzymes are of functional significance to cellular redox homeostasis. Thus, NRX1 may play an important novel role in directly regulating the cellular capacity for H<sub>2</sub>O<sub>2</sub> detoxification, thereby protecting plant cells from oxidative stress triggered by environmental challenges.

## Materials and Methods

Detailed procedures for plant materials and treatments, protein assays, and identification of NRX1 substrates by mass spectrometry are described in SI Materials and Methods.

**ACKNOWLEDGMENTS.** This work was supported by Royal Society University Research Fellowships (UF090321 and UF140600) and Research Grant Rg110495, Wellcome Trust–University of Edinburgh Institutional Strategic Support Fund (ISSF), and European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme Grant No 678511. Further, S.K. was supported by a studentship from the Biotechnology and Biological Sciences Research Council (BBSRC), R.K. by a studentship from the Higher Education Commission Pakistan, G.J.L. by BBSRC Grant BB/D011809/1, T.L.B. by European Union's Seventh Framework Programme 602470, and V.D.-H. and J.-P.R. by the Centre National de la Recherche Scientifique and the Agence Nationale de la Recherche ANR-Blanc Cynthiol 12-BSV6-0011. Liquid chromatography–mass spectrometry (LC–MS) QExactive equipment was purchased by an ISSF award and Wellcome Trust for the Centre for Immunity, Infection and Evolution Award 095831/Z/11/Z.

- Mittler R, et al. (2011) ROS signaling: The new wave? *Trends Plant Sci* 16:300–309.
- Schieber M, Chandel NS (2014) ROS function in redox signaling and oxidative stress. *Curr Biol* 24:R453–R462.
- Petrov VD, Van Breusegem F (2012) Hydrogen peroxide—A central hub for information flow in plant cells. *AoB Plants* 2012:pls014.
- Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7: 405–410.
- Finkel T, Holbrook NJ (2000) Oxidants, oxidative stress and the biology of ageing. *Nature* 408:239–247.
- Day AM, et al. (2012) Inactivation of a peroxiredoxin by hydrogen peroxide is critical for thioredoxin-mediated repair of oxidized proteins and cell survival. *Mol Cell* 45:398–408.
- Meyer Y, Buchanan BB, Vignols F, Reichheld J-P (2009) Thioredoxins and glutaredoxins: Unifying elements in redox biology. *Annu Rev Genet* 43:335–367.
- Foyer CH, Noctor G (2005) Redox homeostasis and antioxidant signaling: A metabolic interface between stress perception and physiological responses. *Plant Cell* 17:1866–1875.
- Perkins A, Nelson KJ, Parsonage D, Poole LB, Karplus PA (2015) Peroxiredoxins: Guardians against oxidative stress and modulators of peroxide signaling. *Trends Biochem Sci* 40:435–445.
- Zamocky M, Furtmüller PG, Obinger C (2008) Evolution of catalases from bacteria to humans. *Antioxid Redox Signal* 10:1527–1548.
- Mittler R, et al. (1999) Transgenic tobacco plants with reduced capability to detoxify reactive oxygen intermediates are hyperresponsive to pathogen infection. *Proc Natl Acad Sci USA* 96:14165–14170.
- Willekens H, et al. (1997) Catalase is a sink for H<sub>2</sub>O<sub>2</sub> and is indispensable for stress defence in C3 plants. *EMBO J* 16:4806–4816.
- Koepke JI, Wood CS, Terlecky LJ, Walton PA, Terlecky SR (2008) Progeric effects of catalase inactivation in human cells. *Toxicol Appl Pharmacol* 232:99–108.
- Terlecky SR, Koepke JI, Walton PA (2006) Peroxisomes and aging. *Biochim Biophys Acta* 1763:1749–1754.
- Marchal C, et al. (2014) NTR/NRX define a new thioredoxin system in the nucleus of *Arabidopsis thaliana* cells. *Mol Plant* 7:30–44.
- Motohashi K, Kondoh A, Stumpp MT, Hisabori T (2001) Comprehensive survey of proteins targeted by chloroplast thioredoxin. *Proc Natl Acad Sci USA* 98:11224–11229.
- Yamazaki D, Motohashi K, Kasama T, Hara Y, Hisabori T (2004) Target proteins of the cytosolic thioredoxins in *Arabidopsis thaliana*. *Plant Cell Physiol* 45:18–27.
- Mhamdi A, et al. (2010) Catalase function in plants: A focus on *Arabidopsis* mutants as stress-mimic models. *J Exp Bot* 61:4197–4220.
- Queval G, et al. (2007) Conditional oxidative stress responses in the *Arabidopsis* photorespiratory mutant *cat2* demonstrate that redox state is a key modulator of daylength-dependent gene expression, and define photoperiod as a crucial factor in the regulation of H<sub>2</sub>O<sub>2</sub>-induced cell death. *Plant J* 52:640–657.
- Chaouch S, et al. (2010) Peroxisomal hydrogen peroxide is coupled to biotic defense responses by ISOCHORISMATE SYNTHASE1 in a daylength-related manner. *Plant Physiol* 153:1692–1705.
- Yuan H-M, Liu W-C, Lu Y-T (2017) CATALASE2 coordinates SA-mediated repression of both auxin accumulation and JA biosynthesis in plant defenses. *Cell Host Microbe* 21:143–155.
- Hayashi T, et al. (2010) Nucleoredoxin negatively regulates Toll-like receptor 4 signaling via recruitment of flightless-1 to myeloid differentiation primary response gene (88). *J Biol Chem* 285:18586–18593.
- Li Y-B, et al. (2016) The thioredoxin GbNRX1 plays a crucial role in homeostasis of apoplastic reactive oxygen species in response to *Verticillium dahliae* infection in cotton. *Plant Physiol* 170:2392–2406.
- Oshima Y, et al. (2008) Plant catalase is imported into peroxisomes by Pex5p but is distinct from typical PTS1 import. *Plant Cell Physiol* 49:671–677.
- Legakis JE, et al. (2002) Peroxisome senescence in human fibroblasts. *Mol Biol Cell* 13: 4243–4255.
- Apanasets O, et al. (2014) PEX5, the shuttling import receptor for peroxisomal matrix proteins, is a redox-sensitive protein. *Traffic* 15:94–103.
- Zou J-J, et al. (2015) *Arabidopsis* CALCIUM-DEPENDENT PROTEIN KINASE8 and CATALASE3 function in abscisic acid-mediated signaling and H<sub>2</sub>O<sub>2</sub> homeostasis in stomatal guard cells under drought stress. *Plant Cell* 27:1445–1460.
- Brown GC (1995) Reversible binding and inhibition of catalase by nitric oxide. *Eur J Biochem* 232:188–191.
- Clark D, Durner J, Navarre DA, Klessig DF (2000) Nitric oxide inhibition of tobacco catalase and ascorbate peroxidase. *Mol Plant Microbe Interact* 13:1380–1384.
- Michelet L, et al. (2013) Down-regulation of catalase activity allows transient accumulation of a hydrogen peroxide signal in *Chlamydomonas reinhardtii*. *Plant Cell Environ* 36:1204–1213.
- Benhar M, Forrester MT, Hess DT, Stamler JS (2008) Regulated protein denitrosylation by cytosolic and mitochondrial thioredoxins. *Science* 320:1050–1054.
- Chae HZ, Chung SJ, Rhee SG (1994) Thioredoxin-dependent peroxide reductase from yeast. *J Biol Chem* 269:27670–27678.
- Kneeshaw S, Gelineau S, Tada Y, Loake GJ, Spoel SH (2014) Selective protein denitrosylation activity of Thioredoxin-h5 modulates plant immunity. *Mol Cell* 56:153–162.
- Montrichard F, et al. (2009) Thioredoxin targets in plants: The first 30 years. *J Proteomics* 72:452–474.
- Pavet V, et al. (2005) Ascorbic acid deficiency activates cell death and disease resistance responses in *Arabidopsis*. *Plant Physiol* 139:1291–1303.
- Provart N, Zhu T (2003) A browser-based functional Classification SuperViewer for *Arabidopsis* genomics. *Curr Comput Mol Biol* 2003:271–272.
- Wildermuth MC, Dewdney J, Wu G, Ausubel FM (2001) Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* 414:562–565.
- Earley KW, et al. (2006) Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J* 45:616–629.
- Clough SJ, Bent AF (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16:735–743.
- Tada Y, et al. (2008) Plant immunity requires conformational changes [corrected] of NPR1 via S-nitrosylation and thioredoxins. *Science* 321:952–956.
- Holmgren A (1979) Thioredoxin catalyzes the reduction of insulin disulfides by dithiothreitol and dihydroliipoamide. *J Biol Chem* 254:9627–9632.
- Bashandy T, et al. (2010) Interplay between the NADP-linked thioredoxin and glutathione systems in *Arabidopsis* auxin signaling. *Plant Cell* 22:376–391.
- Forrester MT, Foster MW, Benhar M, Stamler JS (2009) Detection of protein S-nitrosylation with the biotin-switch technique. *Free Radic Biol Med* 46:119–126.
- Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M (2006) In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nat Protoc* 1:2856–2860.









Fig. S4. NRX1 regulates catalase activity. (A) Protein extracts from untreated WT and *nrx1-1* plants were desalted on NAP-5 columns and protein concentrations equalized. Decomposition of added H<sub>2</sub>O<sub>2</sub> (5 mM) was measured at A<sub>240</sub> and leaf catalase activity (mmol H<sub>2</sub>O<sub>2</sub>·mg·protein<sup>-1</sup>·min<sup>-1</sup>) calculated using the extinction coefficient of 0.036 cm<sup>-2</sup>/mol. Error bars represent SE (n = 3). Asterisks indicate statistically significant differences between samples (Student's t test, P < 0.05). (B) Catalase protein abundance does not differ between WT and *nrx1* mutant lines. Protein was extracted from untreated, MV-treated (5 μM), or Psm ES4326- (5 × 10<sup>5</sup> cells) treated WT, *nrx1-1*, and *cat2 cat3* plants. Analysis was performed by Western blotting using an anti-CAT antibody and an anti-HSP90 antibody as loading control. (C) Protein extract from untreated WT and *nrx1-1* plants was desalted on NAP-5 columns and protein concentrations equalized. Extracts were incubated with either 4 μM recombinant NRX1 or NRX1(C55,58,375,378S) and 0.33 mM DTT. Decomposition of added H<sub>2</sub>O<sub>2</sub> (5 mM) was measured at A<sub>240</sub> and leaf catalase activity (mmol H<sub>2</sub>O<sub>2</sub>·mg·protein<sup>-1</sup>·min<sup>-1</sup>) calculated using the extinction coefficient of 0.036 cm<sup>-2</sup>/mol. Error bars represent SE (n = 3). Asterisks indicate significant differences between WT and mutant recombinant NRX1 proteins (Student's t test, P < 0.05).

## Other Supporting Information Files

[Dataset S1 \(XLSX\)](#)